

LETTER TO EDITOR

Cycling Exercise for Hippocampal Cognitive Function in Older People

Upik Rahmi¹, MD; Hanna Goenawan², PhD; Nova Sylviana², PhD; Iwan Setiawan², PhD

¹Department of Nursing, Faculty of Medicine, Universitas Pendidikan Indonesia, Bandung, Indonesia;

²Department of Biomedical Science, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

Corresponding Author:

Upik Rahmi, MD; Department of Nursing, Faculty of Medicine, Jl. Dr. Setiabudi No. 229 Bandung, Jawa Barat, 40153, Indonesia

Tel/Fax: +622 2004750; Email: upikrahmi@upi.edu

Received: 04 January 2024 Revised: 12 February 2024 Accepted: 14 February 2024

DEAR EDITOR

Several studies have shown the effects of that physical exercise can improve cognitive function.¹ Cycling exercise, as part of aerobic exercise, has a beneficial effect on improving brain health, especially the hippocampus as a cognitive center. Physical exercise increases functional plasticity of the hippocampus by neurogenesis in the Dentate Gyrus, increasing dendritic complexity, spine density, and synaptic plasticity.²

In the elderly, there is a decline in cognitive abilities due to the normal aging process caused by a sedentary lifestyle and cognitive disorders such as dementia and Alzheimer's. Physical exercise can protect the nerves against neurodegenerative disease changes with a structured, individualized exercise program, intensity, and duration to maintain cognitive performance in older adults. Improved cognitive function due to physical exercise influences the expression of the Brain-Derived Neurotrophic Factor (BDNF) gene in the hippocampus, a part of the brain that plays an essential role in learning, memory, and other cognitive functions. BDNF is a neurotrophic factor in brain development, maintenance, and plasticity. BDNF contributes to the growth and development of nerve cells and supports cognitive function and memory development. Regular physical exercise can increase the production and expression of BDNF in the hippocampus. This means physical exercise can help increase the availability of BDNF in the brain, which is stored with various cognitive benefits, including improved memory, learning, and other cognitive functions.³ Mechanisms involved in the relationship between physical exercise and increased BDNF gene expression in the hippocampus are increased blood flow to the brain during exercise, which can increase the supply of oxygen and nutrients to the brain cells, including the cells that produce BDNF. Physical exercise can also stimulate the production of neurotransmitters, such as serotonin, which contributes to increased BDNF.⁴

A study showed that acute cycling exercise with moderate to vigorous intensity for 20 minutes, compared with a control condition of sitting rest can change the hippocampal subfield neurophysiology.⁵ In the hippocampus, cognitive performance increased with 20 minutes of high-intensity cycling exercise as evidenced by salivary alpha-amylase levels in adults. Cycling exercise produced a strong increase in salivary alpha-amylase about six times higher.⁶ Cycling also influences the response speed and the tasks that require attention. Increased hippocampal executive function by cycling outdoors for 16 weeks where function is a cognitive skill for managing oneself and directing behaviours towards certain goals by involving various mental

processes that help a person plan, organize, make decisions, solve problems, control impulses, and monitor action results.⁷ 24 weeks of interactive cycling training had an effect also on the executive function of older adults with mild dementia; in addition, because this study had a combination of cognitive-aerobic training and one aerobic training group, the effectiveness of different interventional components could be identified. The results of this study can be used to recommend physical and mental activity for elderly people with dementia.³

It can be concluded that aerobic exercise, especially cycling both outdoors and indoors with moderate and heavy intensity, can improve the cognitive and executive function in the hippocampus of the elderly and can also be used as a therapy for sufferers of Alzheimer's and dementia.

ACKNOWLEDGMENT

This work was supported by funding from the Research Fund Management Institute (LPDP) from the Indonesian Ministry of Finance, located in the Danadyaksa Cikini Building, Jl. Cikini Raya No. 91 Jakarta Indonesia.

Conflict of Interest: None declared.

Please cite this article as: Rahmi U, Goenawan H, Sylviana N, Setiawan I. Cycling Exercise for Hippocampal Cognitive Function in Older People. *IJCBNM*. 2024;12(3):214-215. doi: 10.30476/IJCBNM.2024.101298.2408.

REFERENCES

- 1 Wang CH, Moreau D, Yang CT, et al. Aerobic exercise modulates transfer and brain signal complexity following cognitive training. *Biological Psychology*. 2019;144:85-98.
- 2 Abdulla SY. Becoming a Supple Leopard: The Ultimate Guide to Resolving Pain, Preventing Injury, and Optimizing Athletic Performance. *The Journal of the Canadian Chiropractic Association*. 2014;58:328.
- 3 Karssemeijer EGA, Bossers WJR, Aaronson JA, et al. The effect of an interactive cycling training on cognitive functioning in older adults with mild dementia: study protocol for a randomized controlled trial. *BMC Geriatrics*. 2017;17:73.
- 4 Jeon YK, Ha CH. The effect of exercise intensity on brain-derived neurotrophic factor and memory in adolescents. *Environmental Health and Preventive Medicine*. 2017;22:27.
- 5 Callow DD, Kommula Y, Stark CEL, Smith JC. Acute cycling exercise and hippocampal subfield function and microstructure in healthy older adults. *Hippocampus*. 2023;33:1123-38.
- 6 Weiss LR, Venezia AC, Smith JC. A single bout of hard RPE-based cycling exercise increases salivary alpha-amylase. *Physiology & Behaviour*. 2019;208:112555.
- 7 Ryu J, Jung JH, Kim J, et al. Outdoor cycling improves clinical symptoms, cognition, and objectively measured physical activity in patients with schizophrenia: A randomized controlled trial. *Journal of Psychiatric Research*. 2020;120:144-53.